

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p504S Antibody / AMACR, Clone: [13H4], Rabbit, Monoclonal NSJ-V8770-20UG

Artikelname	p504S Antibody / AMACR, Clone: [13H4], Rabbit, Monoclonal
Artikelnummer	NSJ-V8770-20UG
Hersteller Artikelnummer	V8770-20UG
Alternativnummer	NSJ-V8770-20UG
Hersteller	NSJ Bioreagents
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC-P, WB
Spezies Reaktivität	Human
Immunogen	Full length human recombinant protein was used as the immunogen for the p504S antibody.
Produktbeschreibung	This antibody recognizes a protein of 44kDa, which is identified as AMACR, also known as p504S. It is an enzyme that is involved in bile acid biosynthesis and beta-oxidation of branched-chain fatty acids. AMACR is essential in lipid metabolism. It is...
Klonalität	Monoclonal
Klon-Bezeichnung	[13H4]
UniProt	Q9UHK6
Reinheit	Protein A affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml, Western blot: 1-2ug/ml
Anwendungsbeschreibung	Optimal dilution of the p504S antibody should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.