

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-MyoD1 Antibody, Clone: [SPM427], Mouse, Monoclonal NSJ-V9052-20UG

Artikelname	Anti-MyoD1 Antibody, Clone: [SPM427], Mouse, Monoclonal
Artikelnummer	NSJ-V9052-20UG
Hersteller Artikelnummer	V9052-20UG
Alternativnummer	NSJ-V9052-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant mouse protein was used as the immunogen for this anti-MyoD1 antibody. The epitope of this mAb maps between amino acid 180-189.
Produktbeschreibung	Recognizes a phosphor-protein of 45kDa, identified as MyoD1. It does not cross react with myogenin, Myf5, or Myf6. Antibody to MyoD1 labels the nuclei of myoblasts in developing muscle tissues. MyoD1 is not detected in normal adult tissue, but is hig...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM427]
UniProt	P15172
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the anti-MyoD1 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9, for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.