

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ornithine Decarboxylase Antibody, Clone: [SPM565], Mouse, Monoclonal NSJ-V9062-100UG

Artikelname	Ornithine Decarboxylase Antibody, Clone: [SPM565], Mouse, Monoclonal
Artikelnummer	NSJ-V9062-100UG
Hersteller Artikelnummer	V9062-100UG
Alternativnummer	NSJ-V9062-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant human protein was used as the immunogen for this ODC-1 antibody.
Produktbeschreibung	Recognizes a 53kDa protein, identified as the Ornithine Decarboxylase (ODC-1). ODC is the initial and rate-limiting enzyme in the biosynthetic pathway of polyamines and is involved in the conversion of ornithine to putrescine. The biological activity...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM565]
UniProt	P11926
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the Ornithine Decarboxylase antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.