

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

ANO1 Antibody / DOG1 / TMEM16A, Clone: [SPM580], Mouse, Monoclonal NSJ-V9068-20UG

Artikelname	ANO1 Antibody / DOG1 / TMEM16A, Clone: [SPM580], Mouse, Monoclonal
Artikelnummer	NSJ-V9068-20UG
Hersteller Artikelnummer	V9068-20UG
Alternativnummer	NSJ-V9068-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant human DOG1 protein was used as the immunogen for this ANO1 antibody.
Produktbeschreibung	Expression of DOG-1/ANO1/TMEM16A protein is elevated in the gastrointestinal stromal tumors (GISTs), c-kit signaling-driven mesenchymal tumors of the GI tract. ANO1 is rarely expressed in other soft tissue tumors, which, due to appearance, may be dif...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM580]
UniProt	Q5XXA6
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the ANO1 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.