

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

ANO1 Antibody / DOG1 / TMEM16A, Clone: [SPM580], Mouse, Monoclonal NSJ-V9068IHC-7ML

Artikelname	ANO1 Antibody / DOG1 / TMEM16A, Clone: [SPM580], Mouse, Monoclonal
Artikelnummer	NSJ-V9068IHC-7ML
Hersteller Artikelnummer	V9068IHC-7ML
Alternativnummer	NSJ-V9068IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant human DOG1 protein was used as the immunogen for this ANO1 antibody.
Produktbeschreibung	Expression of DOG-1/ANO1/TMEM16A protein is elevated in the gastrointestinal stromal tumors (GISTs), c-kit signaling-driven mesenchymal tumors of the GI tract. ANO1 is rarely expressed in other soft tissue tumors, which, due to appearance, may be dif...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM580]
UniProt	Q5XXA6
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	The optimal dilution of the ANO1 antibody for each application should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.