

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Bcl-2 Antibody, Clone: [SPM530], Mouse, Monoclonal NSJ-V9074SAF-100UG

| | |
|--------------------------|---|
| Artikelname | Anti-Bcl-2 Antibody, Clone: [SPM530], Mouse, Monoclonal |
| Artikelnummer | NSJ-V9074SAF-100UG |
| Hersteller Artikelnummer | V9074SAF-100UG |
| Alternativnummer | NSJ-V9074SAF-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FACS, IF, IHC-P, WB |
| Spezies Reaktivität | Human |
| Immunogen | Amino acids 41-54 (GAAPAPGIFSSQPG) from the human protein were used as the immunogen for this anti-Bcl-2 antibody. |
| Produktbeschreibung | This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SPM530] |
| UniProt | P10415 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|--|
| Formulierung | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Flow cytometry: 1-2ug/10 ⁶ cells, Immunofluorescence: 1-2ug/ml, Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | The optimal dilution of the anti-Bcl-2 antibody for each application should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |