

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Bcl-X Antibody, Clone: [SPM165], Mouse, Monoclonal NSJ-V9075-20UG

Artikelname	Anti-Bcl-X Antibody, Clone: [SPM165], Mouse, Monoclonal
Artikelnummer	NSJ-V9075-20UG
Hersteller Artikelnummer	V9075-20UG
Alternativnummer	NSJ-V9075-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Amino acids 3-14 (QSNRELVVDFLS) from the human protein were used as the immunogen for this anti-Bcl-X antibody.
Produktbeschreibung	Recognizes a protein of 27kDa, identified as the Bcl-X protein. This mAb shows no cross-reaction with Bcl-2 or Bax protein. Bcl-X has two isoforms, Bcl-XL (long), a 241 amino acid protein which suppresses cell death. And Bcl-XS (short), a 178 amino a...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM165]
UniProt	<a href="#">Q07817</a>
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the anti-Bcl-X antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.