

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### gp100 Antibody / PMEL17, Clone: [SPM142], Mouse, Monoclonal NSJ-V9078IHC-7ML

Artikelname	gp100 Antibody / PMEL17, Clone: [SPM142], Mouse, Monoclonal
Artikelnummer	NSJ-V9078IHC-7ML
Hersteller Artikelnummer	V9078IHC-7ML
Alternativnummer	NSJ-V9078IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Extract of pigmented melanoma metastases from lymph nodes was used as the immunogen for this gp100 antibody.
Produktbeschreibung	By immunohistochemistry, it specifically recognizes a protein in melanocytes and melanomas. This mAb reacts with junctional and blue nevus cells and variably with fetal and neonatal melanocytes. Intra-dermal nevi, normal adult melanocytes, and non-mel...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM142]
UniProt	<a href="#">P40967</a>
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	The optimal dilution of the gp100 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.