

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### SUMO-2/3 Antibody, Clone: [SPM572], Mouse, Monoclonal NSJ-V9080SAF-100UG

|                          |   |
|--------------------------|---|
| Artikelname              | SUMO-2/3 Antibody, Clone: [SPM572], Mouse, Monoclonal   |
| Artikelnummer            | NSJ-V9080SAF-100UG  |
| Hersteller Artikelnummer | V9080SAF-100UG  |
| Alternativnummer         | NSJ-V9080SAF-100UG  |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IF, IHC-P   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human protein was used as the immunogen for this SUMO-2/3 antibody.   |
| Produktbeschreibung      | This mAb reacts with both SUMO-2 and SUMO-3. The small ubiquitin-related modifier (SUMO) proteins, which include SUMO-1, 2 and 3, belong to the ubiquitin-like protein family. Like ubiquitin, the SUMO proteins are synthesized as precursor proteins tha... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [SPM572]  |
| UniProt                  | <a href="#">P61956</a>  |
| Reinheit                 | Protein G affinity chromatography   |

|                        |   |
|------------------------|---|
| Formulierung           | 1 mg/ml in 1X PBS, BSA free, sodium azide free  |
| Antibody Type          | Primary Antibody  |
| Application Verdünnung | Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 2-4ug/ml for 30 min at RT  |
| Anwendungsbeschreibung | <p>The optimal dilution of the SUMO-2/3 antibody for each application should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p> |