

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

pS2 Antibody / TFF1, Clone: [SPM573], Mouse, Monoclonal NSJ-V9082-20UG

Artikelname	pS2 Antibody / TFF1, Clone: [SPM573], Mouse, Monoclonal
Artikelnummer	NSJ-V9082-20UG
Hersteller Artikelnummer	V9082-20UG
Alternativnummer	NSJ-V9082-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Amino acids 57-84 (CFDDTVRGVPWCFYPNTIDVPPPEEECEF) from the human protein were used as the immunogen for this pS2 antibody.
Produktbeschreibung	It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain t...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM573]
UniProt	P04155
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the pS2 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.