

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

pS2 Antibody / TFF1, Clone: [SPM573], Mouse, Monoclonal NSJ-V9082IHC-7ML

Artikelname	pS2 Antibody / TFF1, Clone: [SPM573], Mouse, Monoclonal
Artikelnummer	NSJ-V9082IHC-7ML
Hersteller Artikelnummer	V9082IHC-7ML
Alternativnummer	NSJ-V9082IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Amino acids 57-84 (CFDDTVRGVPWCFYPNTIDVPPPEEECEF) from the human protein were used as the immunogen for this pS2 antibody.
Produktbeschreibung	It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain t...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM573]
UniProt	P04155
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	The optimal dilution of the pS2 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.