

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Human CD207 (N-6His) EBT-EPT060

Artikelname	Recombinant Human CD207 (N-6His)
Artikelnummer	EBT-EPT060
Hersteller Artikelnummer	EPT060
Alternativnummer	EBT-EPT060-50
Hersteller	ELK Biotechnology
Kategorie	Proteine/Peptide
Produktbeschreibung	Recombinant Human C-type Lectin Domain Family 4 Member K is produced by our Mammalian expression system and the target gene encoding Tyr64-Pro328 is expressed with a 6His tag at the N-terminus....
Molekulargewicht	Molecular weight: 31.5 KDa. Apparent molecular weight: 30-40 KDa, reducing conditions
UniProt	AAH22278.1
Reinheit	Greater than 95% as determined by reducing SDS-PAGE.

Anwendungsbeschreibung	Redissolve: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.. Endotoxin: Less than 0.1 ng/µg (1 EU/µg) as determined by LAL test. Background: Langerin (CD207) is a type II transmembrane glycoprotein which is member K of the C-type lectin domain family. Langerin is used as a marker for Langerhans cells (LCs) which represent the immature dendritic cells in the epidermis. Langerin is necessary and sufficient for Birbeck granule formation. Human langerin sequence contains a 43 aa cytoplasmic domain, a 21 aa transmembrane domain and a 264 aa extracellular domain (ECD) that contains a coiled-coil domain and a single C-type lectin domain. Human langerin shares 68%, 62%, 71% aa identity with mouse, rat and bovine langerin ECD, respectively. Trimerization greatly increases the lectin binding affinity. Langerin internalizes endogenous proteins such as type I procollagen. Internalization by LC is thought to lead to suppression of self reactions. Langerin also mediates endocytosis of non-peptide antigens containing mannose, N-acetyl glucosamine and fucose that are expressed by mycobacteria and fungae
------------------------	--