

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Mouse IL-22 (C-mFc) EBT-EPT148

Artikelname	Recombinant Mouse IL-22 (C-mFc)
Artikelnummer	EBT-EPT148
Hersteller Artikelnummer	EPT148
Alternativnummer	EBT-EPT148-10
Hersteller	ELK Biotechnology
Kategorie	Proteine/Peptide
Produktbeschreibung	Recombinant Mouse Interleukin-22 is produced by our Mammalian expression system and the target gene encoding Leu34-Val179 is expressed with a mFc tag at the C-terminus....
Molekulargewicht	Molecular weight: 43.8 KDa. Apparent molecular weight: 50-65 KDa, reducing conditions
UniProt	Q9JJY9
Reinheit	Greater than 95% as determined by reducing SDS-PAGE.

Anwendungsbeschreibung	Redissolve: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.. Endotoxin: Less than 0.1 ng/µg (1 EU/µg) as determined by LAL test. Background: Interleukin-22 (IL-22) was initially identified as a gene induced by IL-9 in mouse T cells and mast cells. Mouse IL-22 cDNA encodes a 179 amino acid residue protein with a putative 33 amino acid signal peptide that is cleaved to generate a 147 amino acid mature protein that shares approximately 79% and 22% sequence identity with human IL22 and IL10, respectively. IL22 has been shown to activate STAT-1 and STAT-3 in several hepatoma cell lines and up-regulate the production of acute phase proteins. IL-22 is produced by normal mouse T cells upon Con A activation. Mouse IL-22 expression is also induced in various organs upon lipopolysaccharide injection, suggesting that IL-22 may be involved in inflammatory responses. The functional IL-22 receptor complex consists of two receptor subunits, IL-22R (previously an orphan receptor named CRF2-9) and IL-10Rbeta (previously known as CRF2-4), belonging to the class II cytokine receptor family
------------------------	--