

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Nucleoli Marker Antibody / Human Nucleolar Antigen, Clone: [NM95], Unconjugated, Mouse, Monoclonal BYT-ORB2638091

Artikelname	Nucleoli Marker Antibody / Human Nucleolar Antigen, Clone: [NM95], Unconjugated, Mouse, Monoclonal
Artikelnummer	BYT-ORB2638091
Hersteller Artikelnummer	orb2638091
Alternativnummer	BYT-ORB2638091-100
Hersteller	Biorbyt
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, ICC, IF, IHC-P
Spezies Reaktivität	Human
Immunogen	Nuclei of myeloid leukemia biopsy cells were used as the immunogen.
Konjugation	Unconjugated
Produktbeschreibung	This mAb is part of a panel of reagents which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. Clone NM95 antibody rec...
Klonalität	Monoclonal
Klon-Bezeichnung	[NM95]

Puffer	1 mg/ml in 1X PBS, rAlbumin free, sodium azide free
Application Verdünnung	Flow cytometry: 1-2ug/10 ⁶ cells ,Immunofluorescence: 1-2ug/ml,Immunocytochemistry: 1-2ug/ml for 30 min at RT,Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Application Notes: The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the Nucleoli marker antibody to be titrated up or down for optimal performance. 1. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min