

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MUC1 Antibody / Mucin-1 / EMA, Clone: [MUC1/520], Unconjugated, Mouse, Monoclonal BYT-ORB605908

Artikelname	MUC1 Antibody / Mucin-1 / EMA, Clone: [MUC1/520], Unconjugated, Mouse, Monoclonal
Artikelnummer	BYT-ORB605908
Hersteller Artikelnummer	orb605908
Alternativnummer	BYT-ORB605908-100,BYT-ORB605908-20
Hersteller	Biorbyt
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P, WB
Spezies Reaktivität	Human
Immunogen	Human milk-fat globule membranes (HMFGM) was used as the immunogen for the MUC1 antibody.
Konjugation	Unconjugated
Produktbeschreibung	In Western blotting, it recognizes proteins in MW range of 265-400 kDa, identified as different glycoforms of MUC1/Mucin-1/Epithelial Marker Antigen/EMA. The alpha subunit has cell adhesive properties. It can act both as an adhesion and an anti-adhes...
Klonalität	Monoclonal
Klon-Bezeichnung	[MUC1/520]

UniProt	P15941
Puffer	0.2 mg/ml in 1X PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT, Western blot: 1-2ug/ml
Anwendungsbeschreibung	Application Notes: Optimal dilution of the MUC1 antibody should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min