

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Swine anti Horse Ig light chains kappa and lambda (free and bound), Clone: [Polyclonal], Porcine, Monoclonal NMB-SWAHO/L(SD+HD)

Artikelname	Swine anti Horse Ig light chains kappa and lambda (free and bound), Clone: [Polyclonal], Porcine, Monoclonal
Artikelnummer	NMB-SWAHO/L(SD+HD)
Hersteller Artikelnummer	SwAHo/L(SD+HD)
Alternativnummer	NMB-SWAHO/L(SD+HD)
Hersteller	NordicMubio
Wirt	Porcine
Kategorie	Antikörper
Produktbeschreibung	As tested in immunoelectrophoresis and double radial immunodiffusion against pooled normal horse serum and purified IgG fractions and light chain preparations, the antiserum reacts with determinants exposed on the surface (SD) and the Fab portion of ...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]
Puffer	Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled swine ser
Quelle	Purified equine light chains isolated from polyclonal horse IgG. Freund's complete adjuvant is used in the first step of the immunization procedure.

Formel	Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled swine se
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and serially diluted) in 2 mm diameter peripheral wells. In single immunodiffusion, use 1% antiserum in agar gel to establish a standard curve.