

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Swine anti Rabbit IgG (heavy and light chains), Clone: [Polyclonal], Porcine, Monoclonal NMB-SWAR/IGG(H+L)

Artikelname	Swine anti Rabbit IgG (heavy and light chains), Clone: [Polyclonal], Porcine, Monoclonal
Artikelnummer	NMB-SWAR/IGG(H+L)
Hersteller Artikelnummer	SwAR/IgG(H+L)
Alternativnummer	NMB-SWAR/IGG(H+L)
Hersteller	NordicMubio
Wirt	Porcine
Kategorie	Antikörper
Spezies Reaktivität	Rabbit
Konjugation	Unconjugated
Format	Antiserum
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in rabbit serum, a...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]

Puffer	Purified hyperimmune swine IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. No foreign protein added. It is reconstituted by adding 1 ml sterile distilled wa
Quelle	Purified normal IgG isolated from pooled rabbit serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formel	Purified hyperimmune swine IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. No foreign protein added.
Antibody Type	Secondary Antibody
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells.