

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Bov Ig Light Chains Purified, Clone: [IVA285-1], Monoclonal EXB-11-420-C100

|                          |                                                                  |
|--------------------------|------------------------------------------------------------------|
| Artikelname              | Anti-Bov Ig Light Chains Purified, Clone: [IVA285-1], Monoclonal |
| Artikelnummer            | EXB-11-420-C100                                                  |
| Hersteller Artikelnummer | 11-420-C100                                                      |
| Alternativnummer         | EXB-11-420-C100                                                  |
| Hersteller               | EXBIO                                                            |
| Kategorie                | Antikörper                                                       |
| Applikation              | FC, IHC-Fr, WB                                                   |
| Spezies Reaktivität      | Bovine, Sheep                                                    |
| Immunogen                | Fraction of Bovine immunoglobulin.                               |
| Klonalität               | Monoclonal                                                       |
| Konzentration            | 1 mg/ml                                                          |
| Klon-Bezeichnung         | [IVA285-1]                                                       |
| Isotyp                   | Mouse IgG1                                                       |
| Puffer                   | Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide      |
| Lagerung                 | 2°C to 8°C                                                       |
| Target-Kategorie         | Ig Light Chains                                                  |
| Antibody Type            | Monoclonal Antibody                                              |

|                        |                                                                                                                                                                                                                                                                                                                                                                |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Anwendungsbeschreibung | Western blotting: Recommended dilution: 0,5-1 µg/ml, overnight at 4°C. Sample preparation: bovine serum, dilution 1:50 in Laemmli reducing buffer, boiled in water bath for 3 min. Reducing conditions, 12% separating gel. The antibody strongly reacts with bovine IgG light chains, weakly reacts with IgM. Flow cytometry: Recommended dilution: 1-4 µg/ml |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|