

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit IgG anti-Swine IgG (F(ab)2)-HRPO, MinX none DNA-SEC-182633

Artikelname	Rabbit IgG anti-Swine IgG (F(ab)2)-HRPO, MinX none
Artikelnummer	DNA-SEC-182633
Hersteller Artikelnummer	SEC-182633
Alternativnummer	DNA-SEC-182633
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Porcine
Immunogen	Swine IgG F(ab)2 fragment
Konjugation	HRPO
Format	IgG
Spezifität	IgG (F(ab')2)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Anti-Swine IgG F(ab)2 Antibody generated in rabbit recognizes the dimeric Fab portion of the swine IgG molecule. Swine IgG F(ab)2 is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlle...
Klonalität	Polyclonal

Konzentration	10.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Swine IgG, Swine IgG F(ab') ₂ and Swine Serum. No reaction was observed against Swine IgG F(c).
Formulierung	Lyophilized
Formel	20 mM K ₃ PO ₄ , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% Gentamicin
Target-Kategorie	Swine
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Anwendungsbeschreibung	This product has been assayed against 1.0 ug of Swine IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:27,000 to 1:110,000 of the reconstitution concentration is suggested for this product.