

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Donkey IgG anti-Chicken IgG (H+L)-HRPO, MinX Bo,Go,Gp,Hm,Ho,Hu,Ms,Rb,Rt,Sh DNA-SEC-182807

Artikelname	Donkey IgG anti-Chicken IgG (H+L)-HRPO, MinX Bo,Go,Gp,Hm,Ho,Hu,Ms,Rb,Rt,Sh
Artikelnummer	DNA-SEC-182807
Hersteller Artikelnummer	SEC-182807
Alternativnummer	DNA-SEC-182807
Hersteller	dianova
Wirt	Donkey
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Gallus
Immunogen	Chicken IgG whole molecule
Konjugation	HRPO
Format	IgG
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	Bovine,Goat,Guinea pig,Hamster (all),Equine,Human,Mouse,Rabbit,Rat,Sheep

Produktbeschreibung	Anti-Chicken IgG Peroxidase Antibody generated in donkey detects chicken IgY. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, ...
Klonalität	Polyclonal
Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads followed by conjugation and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-peroxidase, anti-Donkey Serum, Chicken IgG and Chicken Serum. No reaction was observed against Bovine, Goat, Guinea Pig, Hamster, Horse, Human, Mouse, Rabbit, Rat and Sheep Serum Proteins. This antibody will react with heavy chains of Chicken IgG and with light chains of most Chicken immunoglobulins.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target-Kategorie	Chicken
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:65,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:5,000
Anwendungsbeschreibung	Anti-Chicken IgG (H&L) is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the end user.