

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Donkey IgG anti-Goat IgG (H+L)-HRPO, MinX Hm,Rb,Rt,Ck,Gp,Ho,Ms DNA-SEC-182886

Artikelname	Donkey IgG anti-Goat IgG (H+L)-HRPO, MinX Hm,Rb,Rt,Ck,Gp,Ho,Ms
Artikelnummer	DNA-SEC-182886
Hersteller Artikelnummer	SEC-182886
Alternativnummer	DNA-SEC-182886
Hersteller	dianova
Wirt	Donkey
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Goat
Immunogen	Anti-Goat IgG (H&L) was produced by repeated immunization with Goat IgG whole molecule in donkey.
Konjugation	HRPO
Format	IgG
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	Hamster (all),Rabbit,Rat,Gallus,Guinea pig,Equine,Mouse
Produktbeschreibung	Anti-Goat IgG Peroxidase antibody generated in donkey detects goat IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as wel...

Klonalität	Polyclonal
Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Donkey Serum, Goat IgG and Goat Serum. No reaction was observed against Chicken, Guinea Pig, Hamster, Horse, Mouse, Rabbit and Rat Serum Proteins.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target-Kategorie	Goat
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:170,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:2,000 - 1:10,000
Anwendungsbeschreibung	Anti-Goat IgG (H&L) is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the end user.