

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Human Lambda light chain-HRPO, MinX none DNA-SEC-183027

Artikelname	Goat IgG anti-Human Lambda light chain-HRPO, MinX none
Artikelnummer	DNA-SEC-183027
Hersteller Artikelnummer	SEC-183027
Alternativnummer	DNA-SEC-183027
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Human
Immunogen	Anti-Human lambda chain was produced by repeated immunization with human lambda light chain in goat.
Konjugation	HRPO
Format	IgG
Spezifität	Lambda (light chain)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	The anti-Human lambda (lambda chain) Antibody detects the lambda chain subunit. Immunoglobulins are heterotetramers composed of 2 immunoglobulin heavy and 2 immunoglobulin light chains. The immunoglobulin light chain is the small polypeptide subunit ...

Klonalität	Polyclonal
Konzentration	1.0 mg/ml
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-Goat Serum. Specificity was confirmed by ELISA minimal cross reactivity against other human heavy or light chain isotypes.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Thimerosal
Target-Kategorie	Human
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Anwendungsbeschreibung	This secondary peroxidase conjugated antibody anti-Human lambda chain is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays. Human lambda chain Antibody peroxidase conjugated has been assayed against 1.0 ug of Human IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:16,000 to 1:64,000 is suggested for this product.