

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit IgG anti-Human IgG (H)-Biotin, MinX none DNA-SEC-183090

Artikelname	Rabbit IgG anti-Human IgG (H)-Biotin, MinX none
Artikelnummer	DNA-SEC-183090
Hersteller Artikelnummer	SEC-183090
Alternativnummer	DNA-SEC-183090
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Human
Immunogen	Human IgG gamma heavy chain
Konjugation	Biotin
Format	IgG
Spezifität	IgG (Fc)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Anti-Human IgG (gamma chain) Biotin generated in rabbit detects human Immunoglobulin G (gamma chain). It is a protein complex composed of four peptide chains - two identical heavy chains and two identical light chains arranged in a Y-shape typical of...
Klonalität	Polyclonal

Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Rabbit Serum, Human IgG and Human Serum. No reaction was observed against other Human heavy or light chain proteins.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target-Kategorie	Human
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Anwendungsbeschreibung	Anti-Human IgG (gamma chain) Biotin has been assayed against 1.0 ug of Human IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:5,000 to 1:35,000 of the reconstitution concentration is suggested for this product.