

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Rabbit IgG (Fc)-unconj., MinX none DNA-SEC-183352

Artikelname	Goat IgG anti-Rabbit IgG (Fc)-unconj., MinX none
Artikelnummer	DNA-SEC-183352
Hersteller Artikelnummer	SEC-183352
Alternativnummer	DNA-SEC-183352
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Rabbit
Immunogen	Anti-Rabbit IgG was produced by repeated immunization with rabbit IgG f(c) fragment in goat
Konjugation	Unconjugated
Format	IgG
Spezifität	IgG (Fc)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Anti-Rabbit IgG F(c) generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of rabbit IgG and o...

Klonalität	Polyclonal
Konzentration	2.26 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rabbit IgG, Rabbit IgG F(c) and Rabbit Serum. No reaction was observed against Rabbit IgG F(ab') ₂ .
Formulierung	Liquid (sterile filtered)
Formel	20 mM K ₃ PO ₄ , 150 mM NaCl, pH 7.2, sterile filtered, 0.01% NaN ₃
Target-Kategorie	Rabbit
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Anwendungsbeschreibung	Antibody Anti-Rabbit IgG F(c) has been tested by ELISA, dot blot, and western blot and is suitable for immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring lot-to-lot consistency.