

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Sheep IgG anti-Rabbit IgG (H+L)-HRPO, MinX Bo,Ck,Go,Gp,Ho,Hu,Ms,Rt,Sh DNA-SEC-183412

Artikelname	Sheep IgG anti-Rabbit IgG (H+L)-HRPO, MinX Bo,Ck,Go,Gp,Ho,Hu,Ms,Rt,Sh
Artikelnummer	DNA-SEC-183412
Hersteller Artikelnummer	SEC-183412
Alternativnummer	DNA-SEC-183412
Hersteller	dianova
Wirt	Sheep
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Rabbit
Immunogen	Rabbit IgG whole molecule
Konjugation	HRPO
Format	IgG
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	Bovine,Gallus,Goat,Guinea pig,Equine,Human,Mouse,Rat,Sheep

Produktbeschreibung	Anti-Rabbit IgG (H&L) generated in sheep detects rabbit Immunoglobulin G. Both the Heavy and Light chains of the antibody molecule are present. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found i...
Klonalität	Polyclonal
Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Sheep Serum, Rabbit IgG and Rabbit Serum. No reaction was observed against or Bovine, Chicken, Goat, Guinea Pig, Horse, Human, Mouse, Rat and Sheep Serum Proteins.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target-Kategorie	Rabbit
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Anwendungsbeschreibung	This product has been assayed against 1.0 µg of Rabbit IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:8,000 to 1:40,000 of the reconstitution concentration is suggested for this product.