

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Rat IgG+IgM+IgA (H+L)-HRPO, MinX none DNA-SEC-183446

Artikelname	Goat IgG anti-Rat IgG+IgM+IgA (H+L)-HRPO, MinX none
Artikelnummer	DNA-SEC-183446
Hersteller Artikelnummer	SEC-183446
Alternativnummer	DNA-SEC-183446
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Rat
Immunogen	Rat IgG, IgA and IgM whole molecules
Konjugation	HRPO
Format	IgG
Spezifität	IgG+IgM+IgA (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Rat IgG IgA IgM (H&L) Antibody detects immunoglobulin G, A, and M from Rat. Immunoglobulin G binds to antigens and can neutralize or opsonize targets, and are predominantly involved in the secondary immune response. Immunoglobulin A (IgA) is an antib...
Klonalität	Polyclonal

Konzentration	2.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	Anti-Rat IgG IgA IgM (H&L) Antibody, Peroxidase Conjugated, was prepared from polyspecific antiserum by immunoaffinity chromatography using Rat antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-Goat Serum. This reagent is suitable for the detection of all Rat immunoglobulin isotypes and chain combinations.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Thimerosal
Target-Kategorie	Rat
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:100,000, Immunohistochemistry Dilution: 1:500-1:1,500, Western Blot Dilution: 1:500-1:3,000
Anwendungsbeschreibung	Anti-Rat IgG IgA IgM Antibody is suitable for immunoassays where reactivity to different classes of rat immunoglobulins is desired. Antibody has been tested by ELISA, western blot immunoblot, and immunohistochemistry. Optimal concentrations in immunoassays should be determined by the researcher.