

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Rat IgM (μ)-Alk. Phos., MinX none DNA-SEC-183470

Artikelname	Goat IgG anti-Rat IgM (μ)-Alk. Phos., MinX none
Artikelnummer	DNA-SEC-183470
Hersteller Artikelnummer	SEC-183470
Alternativnummer	DNA-SEC-183470
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Rat
Immunogen	Rat IgM whole molecule
Konjugation	Alk. Phos.
Format	IgG
Spezifität	IgM (μ)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Anti-Rat IgM antibody generated in goat specifically detects rat IgM mu heavy chain. Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly produced in the spleen. Fo...
Klonalität	Polyclonal

Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgM coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Goat Serum, Rat IgM and Rat Serum. No reaction was observed against other rat heavy or light chain proteins.
Formulierung	Liquid (sterile filtered)
Formel	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3
Target-Kategorie	Rat
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:10,000 - 1:30,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Anwendungsbeschreibung	This product has been assayed against 1.0 ug of Rat IgM in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:500 to 1:2,000 is suggested for this product.