

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit IgG anti-Sheep IgG (F(ab)2)-Alk. Phos., MinX none DNA-SEC-183557

Artikelname	Rabbit IgG anti-Sheep IgG (F(ab)2)-Alk. Phos., MinX none
Artikelnummer	DNA-SEC-183557
Hersteller Artikelnummer	SEC-183557
Alternativnummer	DNA-SEC-183557
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Sheep
Immunogen	Sheep IgG F(ab)2 fragment
Konjugation	Alk. Phos.
Format	IgG
Spezifität	IgG (F(ab')2)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Anti-Sheep IgG F(ab)2 Antibody generated in rabbit recognizes the dimeric Fab portion of the sheep IgG molecule. Sheep IgG F(ab)2 is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlle...
Klonalität	Polyclonal

Konzentration	0.6 mg/mL
Isotyp	Ig
Puffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Sheep IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Rabbit Serum, Sheep IgG, Sheep IgG F(ab') ₂ and Sheep Serum. No reaction was observed against Sheep IgG F(c).
Formulierung	Liquid (sterile filtered)
Formel	50 mM TrisHCl, 150 mM NaCl, 1 mM MgCl, 0.1 mM ZnCl, 50% (v/v) Glycerol, pH 8.0, sterile filtered, 0.01% NaN ₃
Target-Kategorie	Sheep
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Anwendungsbeschreibung	This product has been assayed against 1.0 µg of Sheep IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:500 to 1:2,000 of the reconstitution concentration is suggested for this product.