

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Donkey IgG anti-Sheep IgG (H+L)-Alk. Phos., MinX Ck,Gp,Hm,Ho,Hu,Ms,Rb,Rt DNA-SEC-183571

Artikelname	Donkey IgG anti-Sheep IgG (H+L)-Alk. Phos., MinX Ck,Gp,Hm,Ho,Hu,Ms,Rb,Rt
Artikelnummer	DNA-SEC-183571
Hersteller Artikelnummer	SEC-183571
Alternativnummer	DNA-SEC-183571
Hersteller	dianova
Wirt	Donkey
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Sheep
Immunogen	Sheep IgG whole molecule
Konjugation	Alk. Phos.
Format	IgG
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	Gallus,Guinea pig,Hamster (all),Equine,Human,Mouse,Rabbit,Rat

Produktbeschreibung	Anti-Sheep IgG (H&L) generated in donkey detects sheep Immunoglobulin G. Both the Heavy and Light chains of the antibody molecule are present. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in...
Klonalität	Polyclonal
Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Sheep IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Donkey Serum, Sheep IgG and Sheep Serum. No reaction was observed against Chicken, Guinea Pig, Hamster, Horse, Human, Mouse, Rabbit and Rat Serum Proteins.
Formulierung	Liquid (sterile filtered)
Formel	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3
Target-Kategorie	Sheep
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Anwendungsbeschreibung	This product has been assayed against 1.0 ug of Sheep IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:3,000 to 1:14,000 of the reconstitution concentration is suggested for this product.