

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat Fab anti-Human IgG (H+L)-HRPO, MinX none DNA-SEC-183949

Artikelname	Goat Fab anti-Human IgG (H+L)-HRPO, MinX none
Artikelnummer	DNA-SEC-183949
Hersteller Artikelnummer	SEC-183949
Alternativnummer	DNA-SEC-183949
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Human
Immunogen	Human IgG whole molecule
Konjugation	HRPO
Format	Fab
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Fab Anti-Human IgG (H&L) Peroxidase Antibody generated in goat detects immunoglobulin g from human, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum immu...
Klonalität	Polyclonal

Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-Goat Serum. No reaction was observed against anti-Papain or anti-Goat IgG F(c).
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target-Kategorie	Human
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:10,000 - 50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:5,000
Anwendungsbeschreibung	Fab Anti-Human IgG (H&L) Peroxidase Antibody is suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.