

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Hamster IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-GAHA/IGG(H+L)

Article Name	Goat anti Hamster IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GAHA/IGG(H+L)
Supplier Catalog Number	GAHa/IgG(H+L)
Alternative Catalog Number	NMB-GAHA/IGG(H+L)
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Hamster (all)
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Human, Porcine
Product Description	The reactivity of the antiserum is directed to the Fc and Fab subunits of IgG. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. In immunoelectrophoresis against hamster serum a single characteristic pr...
Clonality	Monoclonal
Clone Designation	[Polyclonal]

Buffer	Delipidated, heat inactivated lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum
Source	Purified normal IgG isolated from pooled Syrian hamster serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formula	Delipidated, heat inactivated lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum
Antibody Type	Secondary Antibody
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells.