

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Rat IgG1 IgG2a IgG2b IgG2c IgA IgM (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal NMB-GARA/IG(FC)/TRITC

Article Name	Goat anti Rat IgG1 IgG2a IgG2b IgG2c IgA IgM (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GARA/IG(FC)/TRITC
Supplier Catalog Number	GARa/Ig(Fc)/TRITC
Alternative Catalog Number	NMB-GARA/IG(FC)/TRITC
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Rat
Conjugation	TRITC
Format	IgG
Target Specificity	IgG+IgM+IgA (Fc)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is directed to the Fc subunits of the major isotypes of the rat immunoglobulin system. No reaction is obtained with purified IgG/Fab or any non-Ig protein of rat serum, as tested by immunoelectrophoresis and double rad...
Clonality	Monoclonal
Clone Designation	[Polyclonal]

Buffer	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water.
Source	Purified polyclonal rat IgG, and homogenous IgA and IgM isolated from rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Form	TRITC
Formula	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Antibody Type	Secondary Antibody
Application Notes	ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.