

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mouse anti CD65s, IgM, Clone: [VIM2], Monoclonal NMB-GM-4101

Article Name	Mouse anti CD65s, IgM, Clone: [VIM2], Monoclonal
Biozol Catalog Number	NMB-GM-4101
Supplier Catalog Number	GM-4101
Alternative Catalog Number	NMB-GM-4101
Manufacturer	NordicMubio
Host	Mouse
Category	Antikörper
Application	FC, IF
Species Reactivity	Human
Product Description	The epitope recognized by antibody VIM2 is expressed by virtually all myeloid cells including normal and malignant granulocytes and monocytes. In normal myelopoiesis VIM2 can first be detected after the late CFU-GM stage. In acute myeloid leukemias (...)
Clonality	Monoclonal
Clone Designation	[VIM2]
Isotype	IgM
Buffer	PBS pH 7.2, 1% BSA, 0.05% NaN3
Formula	PBS pH 7.2, 1% BSA, 0.05% NaN3

Application Notes

Staining Procedure Direct Immunofluorescence (Staining Procedure)
Nordic-MUBio fluorochrome labeled antibodies are designed for use with either whole blood or isolated mononuclear cell (MNC) preparations. Proposed staining procedure for whole blood in short: - For each sample add 50 µl of EDTA anti-coagulated blood to a 3-5 ml tube - Add 20 µl of the appropriate Nordic-MUBio monoclonal antibody conjugate - Incubate the tube for 15 minutes at 4°C or at room temperature in the dark - Add 100 µl NM-LYSE (Cat.No. GAS-003) to each tube and incubate for 10 minutes at room temperature - Add 3-4 ml of distilled water and vortex, incubate for 5-10 minutes at room temperature - Centrifuge tube for 5 minutes at 300 g - Aspirate supernatant and resuspend pellet in 0.3 ml of sheath fluid - Analyze immediately or store samples at 2-8°C in the dark and analyze within 24 hours For "No-Wash protocol please refer to www.nordicmubio.com Proposed staining procedure for MNC in short: - Carefully add 20 µl antibody conjugate and 50-100 µl MNC to the bottom of a tube - Vortex at low speed for 1-2 seconds - Incubate for 15-30 minutes at 2-8°C or at room temperature - Centrifuge tubes for 5 minutes at 300 g - Remove supernatant, resuspend cells in 2-5 ml of phosphate buffered saline (PBS) and centrifuge cells again for 5 minutes at 300 g - Remove supernatant and resuspend cells in sheath fluid for immediate analysis or resuspend cells in 0.5 ml 1 % formaldehyde and store them at 2-8°C in the dark. Analyze fixed cells within 24 hours

Indirect Immunofluorescence (Staining Procedure) - Mix 20 µl Nordic-MUBio purified antibody with 50 µl whole blood or MNC suspension - Incubate for 15 minutes at 2-8°C - Wash cells with phosphate buffered saline (PBS) - Add to cell pellet 20 µl of affinity purified, fluorochrome labeled F(ab)₂ anti mouse Ig antibodies - Incubate for 15 minutes at 2-8°C - Wash cells with phosphate buffered saline (PBS) or proceed as described for direct staining