

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mouse anti Myeloperoxidase-C2 (MPO-C2), conjugated to PE, IgG1, Clone: [8E6], Monoclonal NMB-GM-4193

Article Name	Mouse anti Myeloperoxidase-C2 (MPO-C2), conjugated to PE, IgG1, Clone: [8E6], Monoclonal
Biozol Catalog Number	NMB-GM-4193
Supplier Catalog Number	GM-4193
Alternative Catalog Number	NMB-GM-4193
Manufacturer	NordicMubio
Host	Mouse
Category	Antikörper
Application	FC
Species Reactivity	Human
Conjugation	PE
Product Description	Myeloperoxidase (MPO) is a glycoprotein present in the azurophil (primary) granules of myeloid cells, which appears in the myeloblast stage of myeloid cell differentiation. MPO is the most common functional protein of myeloid cells and is involved in ...
Clonality	Monoclonal
Clone Designation	[8E6]
Isotype	IgG1
UniProt	P05164

Buffer	2 ml of PE-conjugated anti Myeloperoxidase-C2 (clone 8E6) in PBS pH 7.2, 1% BSA, and 0.05% NaN ₃ , approximately 100 tests.
Purity	Purified by Chromatography
Form	PE
Formula	PBS pH 7.2, 1% BSA, 0.05% NaN ₃
Application Notes	Permeabilization and Staining Procedure - In combination with our Permeabilization Kit FIX&PERM (Cat. No. GAS-002) intracellular MPO can be easily stained in cell suspensions. - For each sample to be analyzed add 50 µl of whole blood, bone marrow or mononuclear cell suspension in a 5 ml tube - Add 100 µl of Reagent A (Fixation Medium, stored and used at room temperature) - Incubate for 15 minutes at room temperature - Add 5 ml phosphate buffered saline and centrifuge cells for 5 minutes at 300 g - Remove supernatant and add to cell pellet 100 µl Reagent B (Permeabilization Medium) and 20 µl of the MPO-C2 monoclonal antibody conjugate - Vortex at low speed for 1-2 seconds - Incubate for 15 minutes at room temperature - Wash cells with phosphate buffered saline as described above - Remove supernatant and resuspend cells in sheath fluid for immediate analysis or resuspend cells in 0.5 ml 1.0 % formaldehyde and store them at 2-8C in the dark. Analyze fixed cells within 24 hours.