

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Horse IgM (Fc specific), Clone: [Polyclonal], Monoclonal NMB-RAHO/IGM(FC)

Article Name	Rabbit anti Horse IgM (Fc specific), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-RAHO/IGM(FC)
Supplier Catalog Number	RAHo/IgM(Fc)
Alternative Catalog Number	NMB-RAHO/IGM(FC)
Manufacturer	NordicMubio
Host	Rabbit
Category	Antikörper
Species Reactivity	Equine
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgM (μ)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is restricted to the Fc part of the IgM molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal horse serum a single precipitin line is obtained which shows ...
Clonality	Monoclonal
Clone Designation	[Polyclonal]

Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilized
Source	Highly purified normal IgM isolated from pooled horse serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formula	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added.
Antibody Type	Secondary Antibody
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 µl serum, plasma or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and serially diluted) in 2 mm diameter peripheral wells. In single radial immunodiffusion and electroimmunodiffusion use 0.5 to 1.0 percent antiserum in the agar gel.