

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Human Factor VIII related antigen (VWD), Clone: [Polyclonal], Monoclonal NMB-RAHU/FVIII VWD

Article Name	Rabbit anti Human Factor VIII related antigen (VWD), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-RAHU/FVIII VWD
Supplier Catalog Number	RAHu/FVIII VWD
Alternative Catalog Number	NMB-RAHU/FVIII VWD
Manufacturer	NordicMubio
Host	Rabbit
Category	Antikörper
Product Description	The defined antibody reactivity is restricted to VWF. In immunoelectrophoresis, bi-dimensional electrophoresis and radial immunodiffusion (Ouchterlony) against normal plasma, a single precipitin line is obtained which shows a reaction of identity wit...
Clonality	Monoclonal
Clone Designation	[Polyclonal]
UniProt	P04275
Buffer	Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. IgG protein concentration in the antiserum is 10 mg/ml. No foreign proteins added. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water.

Source	Human FVIII procoagulant activity (FVIII:C) is carried by a polypeptide non-covalently bound to a large carrier molecule (unit molecular weight 250 kD) known as the Von Willebrand factor (VWF). FVIII/VWF exists in plasma as a series of polymers with molecu
Formula	Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. IgG protein concentration in the antiserum is 10 mg/ml. No foreign proteins added.
Application Notes	Precipitation assays. The antiserum concentration required in the gel is normally between 1 and 2%. In immunologic determinations of FVIII/VWF, plasma samples and all assay components must contain EDTA to stabilize the proteins The antiserum is standardized for use in the electroimmunodiffusion (EID, Laurell) test procedure for the quantitative determination of FVIII/VWF as described in the Nordic Recommended Working Procedure.