

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

ODC-1 Antibody [ODC1/485], Unconjugated, Mouse, Monoclonal PRS-33-182

Article Name	ODC-1 Antibody [ODC1/485], Unconjugated, Mouse, Monoclonal
Biozol Catalog Number	PRS-33-182
Supplier Catalog Number	33-182
Alternative Catalog Number	PRS-33-182-100
Manufacturer	ProSci
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC, WB
Species Reactivity	Human, Rat
Immunogen	Recombinant human ODC-1 protein was used as the immunogen for this antibody.
Conjugation	Unconjugated
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[ODC1/485]
NCBI	4953
UniProt	P11926
Buffer	PBS with 0.1 mg/ml BSA and 0.05% sodium azide
Form	Liquid

Application Dilute	Optimal dilutions for each application to be determined by the researcher
Application Notes	<p>Flow Cytometry: 0.5-1 ug/million cells IF: 0.5-1 ug/ml WB: 0.5-1 ug/ml IHC (FFPE): 0.5-1 ug/ml for 30 minutes at RT (1) Prediluted format : incubate for 30 min at RT (2) The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the antibody to be titered up or down for optimal performance.</p> <p>1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.</p> <p>2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>