

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

NAIP Peptide PRS-42-537P

Article Name	NAIP Peptide
Biozol Catalog Number	PRS-42-537P
Supplier Catalog Number	42-537P
Alternative Catalog Number	PRS-42-537P-0.1
Manufacturer	ProSci
Category	Proteine/Peptide
Application	Blocking
NCBI	4671
UniProt	Q13075
Buffer	NAIP peptide is supplied as 100 µg of dried peptide. When the peptide is reconstituted in 200 µL water, the concentration will be 0.5 mg/mL.
Form	Lyophilized

Application Notes

NAIP Peptide is used for blocking the activity of NAIP antibody (catalog number 42-537). The NAIP antibody ships at 0.5 mg/mL and the NAIP Peptide ships as a 100 µg pellet. When peptide is reconstituted in 200 µL water, the concentration will be 0.5 mg/mL, identical to that of the antibody. The best starting ratio would be 1:1 (which means molar excess of peptide relative to antibody when identical volumes are mixed). Mix equal volume of peptide and antibody at the required dilution and leave at ambient temperature for 20 minutes. Then incubate and develop in parallel two blots: one with the antibody and one with the antibody which has been pre-absorbed with the provided peptide.