

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

SOX10(SOX10/991 + SOX10/1074), CF640R conjugate, 0.1mg/mL, Clone: [SOX10/991 SOX10/1074], Mouse, Monoclonal BOT-BNC401074-500

| | |
|----------------------------|---|
| Article Name | SOX10(SOX10/991 + SOX10/1074), CF640R conjugate, 0.1mg/mL, Clone: [SOX10/991 SOX10/1074], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC401074-500 |
| Supplier Catalog Number | BNC401074-500 |
| Alternative Catalog Number | BOT-BNC401074-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment (aa 115-269) of human SOX10 protein with hexa-histidine tag |
| Conjugation | CF640R |
| Product Description | Recognizes a protein of ~55 kDa, identified as SOX10. This MAb is highly specific and does not cross-react with other members of the SOX-family. SOX genes comprise a family of genes that are related to the mammalian sex-determining gene SRY. These ge... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [SOX10/991 SOX10/1074] |

| | |
|-------------------|---|
| Molecular Weight | 49-58 kDa |
| UniProt | P56693 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user |