

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin 6 (KRT6) (Hyperproliferation-Related Keratin) (KRT6/1702), CF640R conjugate, 0.1mg/mL, Clone: [KRT6/1702], Mouse, Monoclonal BOT-BNC401702-100

| | |
|----------------------------|--|
| Article Name | Cytokeratin 6 (KRT6) (Hyperproliferation-Related Keratin) (KRT6/1702), CF640R conjugate, 0.1mg/mL, Clone: [KRT6/1702], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC401702-100 |
| Supplier Catalog Number | BNC401702-100 |
| Alternative Catalog Number | BOT-BNC401702-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human, Mouse |
| Immunogen | Recombinant human full-length keratin 6 protein |
| Conjugation | CF640R |
| Product Description | This MAb recognizes a protein of 56 kDa, identified as cytokeratin 6 (CK6). In humans, multiple isoforms of Cytokeratin 6 (6A-6F), encoded by several highly homologous genes, have distinct tissue expression patterns, and Cytokeratin 6A is the dominant... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [KRT6/1702] |

| | |
|-------------------|--|
| Molecular Weight | 56 kDa |
| UniProt | P02538 , P48668 , P04259 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 0.1-0.2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |