

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Parathyroid Hormone (PTH) (N-Terminal) (PTH/1717R), CF640R conjugate, 0.1mg/mL, Clone: [PTH/1717R], Rabbit, Monoclonal BOT-BNC401717-100

| | |
|----------------------------|---|
| Article Name | Parathyroid Hormone (PTH) (N-Terminal) (PTH/1717R), CF640R conjugate, 0.1mg/mL, Clone: [PTH/1717R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNC401717-100 |
| Supplier Catalog Number | BNC401717-100 |
| Alternative Catalog Number | BOT-BNC401717-100-100UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant human PTH protein N-terminal fragment (exact sequence is proprietary) |
| Conjugation | CF640R |
| Product Description | Epitope of this MAb maps in the N-terminus of PTH, a hormone produced by the parathyroid gland that regulates the concentration of calcium and phosphorus in extracellular fluid. This hormone elevates blood calcium levels by dissolving the salts in bo... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [PTH/1717R] |

| | |
|-------------------|--|
| Molecular Weight | 9 kDa |
| UniProt | P01270 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Immunofluorescence 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |