

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) (CFTR/1775R), CF640R conjugate, 0.1mg/mL, Clone: [CFTR/1775R], Rabbit, Monoclonal BOT-BNC401775-500

| | |
|----------------------------|---|
| Article Name | CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) (CFTR/1775R), CF640R conjugate, 0.1mg/mL, Clone: [CFTR/1775R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNC401775-500 |
| Supplier Catalog Number | BNC401775-500 |
| Alternative Catalog Number | BOT-BNC401775-500-500UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant human CFTR fragment (aa258-385) (exact sequence is proprietary) |
| Conjugation | CF640R |
| Product Description | This antibody recognizes a protein of 165-170 kDa, identified as cystic fibrosis transmembrane conductance regulator (CFTR). CFTR is composed of two membrane-spanning domains (MSD), two nucleotide-binding domains (NBD), and an R domain. It is structu... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |

| | |
|-------------------|--|
| Clone Designation | [CFTR/1775R] |
| Molecular Weight | 165-170 kDa |
| UniProt | P13569 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user |