

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human IgG (Immunoglobulin Gamma Heavy Chain) (B-Cell Marker) (rIG266), CF640R conjugate, 0.1mg/mL, Clone: [rIG266], Mouse, Monoclonal BOT-BNC401789-100

| | |
|----------------------------|---|
| Article Name | Human IgG (Immunoglobulin Gamma Heavy Chain) (B-Cell Marker) (rIG266), CF640R conjugate, 0.1mg/mL, Clone: [rIG266], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC401789-100 |
| Supplier Catalog Number | BNC401789-100 |
| Alternative Catalog Number | BOT-BNC401789-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human IGHG protein |
| Conjugation | CF640R |
| Product Description | Recognizes a protein of 75 kDa, identified as gamma heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), mu (IgM), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. This MAb is ... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |

| | |
|-------------------|---|
| Clone Designation | [rIG266] |
| Molecular Weight | 75 kDa |
| UniProt | P01857 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 min at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min |