

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Ferritin, Light Chain (FTL) (Microglia Marker) (FTL/2338R), CF647 conjugate, 0.1mg/mL, Clone: [FTL/2338R], Rabbit, Monoclonal BOT-BNC472338-100

Article Name	Ferritin, Light Chain (FTL) (Microglia Marker) (FTL/2338R), CF647 conjugate, 0.1mg/mL, Clone: [FTL/2338R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNC472338-100
Supplier Catalog Number	BNC472338-100
Alternative Catalog Number	BOT-BNC472338-100-100UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human
Immunogen	Recombinant human FTL protein fragment (around aa 38-165) (exact sequence is proprietary)
Conjugation	CF647
Product Description	Mammalian ferritins consist of 24 subunits made up of 2 types of polypeptide chains, ferritin heavy chain and ferritin light chain. Ferritin heavy chains catalyze the first step in iron storage, the oxidation of Fe (II), whereas ferritin light chains...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[FTL/2338R]

Molecular Weight	19-25 kDa
UniProt	P02792
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Western Blotting 0.5-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user