

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **EGFR (Epidermal Growth Factor Receptor) (GFR/2341), CF647 conjugate, 0.1mg/mL, Clone: [GFR/2341], Mouse, Monoclonal BOT-BNC472341-100**

|                            |   |
|----------------------------|---|
| Article Name               | EGFR (Epidermal Growth Factor Receptor) (GFR/2341), CF647 conjugate, 0.1mg/mL, Clone: [GFR/2341], Mouse, Monoclonal   |
| Biozol Catalog Number      | BOT-BNC472341-100   |
| Supplier Catalog Number    | BNC472341-100   |
| Alternative Catalog Number | BOT-BNC472341-100-100UL   |
| Manufacturer               | Biotium   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | IHC, WB   |
| Species Reactivity         | Human   |
| Immunogen                  | Recombinant fragment of human EGFR protein (around aa 344-492) (exact sequence is proprietary)  |
| Conjugation                | CF647   |
| Product Description        | This MAb recognizes a protein of 170 kDa, identified as EGFR. EGFR is type I receptor tyrosine kinase with sequence homology to erbB-1, -2, -3 -4 or HER-1, -2, -3 -4. It binds to Epidermal Growth Factor (EGF), Transforming Growth Factor-a (TGF-a), He... |
| Clonality                  | Monoclonal  |
| Concentration              | 0.1 mg/mL   |
| Clone Designation          | [GFR/2341]  |

|                   |   |
|-------------------|---|
| Molecular Weight  | ~170 kDa (wild type) and ~145 kDa (vIII variant)  |
| UniProt           | <a href="#">P00533</a>  |
| Buffer            | PBS, 0.1% BSA, 0.05% azide  |
| Source            | Animal  |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |