

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **p63 (Squamous, Basal & Myoepithelial Cell Marker) (TP63/2428), CF647 conjugate, 0.1mg/mL, Clone: [TP63/2428], Mouse, Monoclonal BOT-BNC472428-500**

|                            |                                                                                                                                                                                                                                                               |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name               | p63 (Squamous, Basal & Myoepithelial Cell Marker) (TP63/2428), CF647 conjugate, 0.1mg/mL, Clone: [TP63/2428], Mouse, Monoclonal                                                                                                                               |
| Biozol Catalog Number      | BOT-BNC472428-500                                                                                                                                                                                                                                             |
| Supplier Catalog Number    | BNC472428-500                                                                                                                                                                                                                                                 |
| Alternative Catalog Number | BOT-BNC472428-500-500UL                                                                                                                                                                                                                                       |
| Manufacturer               | Biotium                                                                                                                                                                                                                                                       |
| Host                       | Mouse                                                                                                                                                                                                                                                         |
| Category                   | Antikörper                                                                                                                                                                                                                                                    |
| Application                | FC, IHC                                                                                                                                                                                                                                                       |
| Species Reactivity         | Human                                                                                                                                                                                                                                                         |
| Immunogen                  | Recombinant full-length human p63 protein                                                                                                                                                                                                                     |
| Conjugation                | CF647                                                                                                                                                                                                                                                         |
| Product Description        | p63 is a homolog of the tumor suppressor p53. It is identified in basal cells in the epithelial layers of a variety of tissues, including epidermis, cervix, urothelium, breast and prostate. p63 was detected in nuclei of the basal epithelium in normal... |
| Clonality                  | Monoclonal                                                                                                                                                                                                                                                    |
| Concentration              | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Clone Designation          | [TP63/2428]                                                                                                                                                                                                                                                   |

|                   |                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|-------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molecular Weight  | 63 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| UniProt           | <a href="#">Q9H3D4</a>                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Buffer            | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Source            | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |