

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD106 / VCAM1(1.4C3), CF568 conjugate, 0.1mg/mL, Clone: [1.4C3], Mouse, Monoclonal BOT-BNC680149-100

| | |
|----------------------------|---|
| Article Name | CD106 / VCAM1(1.4C3), CF568 conjugate, 0.1mg/mL, Clone: [1.4C3], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC680149-100 |
| Supplier Catalog Number | BNC680149-100 |
| Alternative Catalog Number | BOT-BNC680149-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IHC, IP, WB |
| Species Reactivity | Human |
| Immunogen | Stimulated human umbilical vein endothelial cells (HUVEC) |
| Conjugation | CF568 |
| Product Description | CD106 (also known as vascular cell adhesion molecule-1, or VCAM-1) is a member of the Ig superfamily of adhesion molecules. It is expressed at high levels on cytokine stimulated vascular endothelial cells, and at minimal levels on un-stimulated endot... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [1.4C3] |

| | |
|-------------------|---|
| Molecular Weight | ~110 kDa |
| UniProt | P19320 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |