

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin 17(E3), CF568 conjugate, 0.1mg/mL, Mouse BOT-BNC680158-100

| | |
|----------------------------|---|
| Article Name | Cytokeratin 17(E3), CF568 conjugate, 0.1mg/mL, Mouse |
| Biozol Catalog Number | BOT-BNC680158-100 |
| Supplier Catalog Number | BNC680158-100 |
| Alternative Catalog Number | BOT-BNC680158-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC, WB |
| Species Reactivity | Bovine, Goat, Human, Porcine, Rat |
| Immunogen | The cytoskeletal fraction of rat colon epithelium |
| Conjugation | CF568 |
| Product Description | Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excellent tool to distinguish myoepithelial cells from luminal epithelium of various glands such as ... |
| Concentration | 0.1 mg/mL |
| Molecular Weight | 46 kDa |
| UniProt | Q04695 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |

| Source | Animal |
|-------------------|---|
| Application Notes | Immunohistology frozen & formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |