

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

ZAP70 (Chronic Lymphocytic Leukemia Marker) (ZAP70/2046), 0.2mg/mL, Clone: [ZAP70/2046], Mouse, Monoclonal BOT-BNUB2046-100

Article Name	ZAP70 (Chronic Lymphocytic Leukemia Marker) (ZAP70/2046), 0.2mg/mL, Clone: [ZAP70/2046], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB2046-100
Supplier Catalog Number	BNUB2046-100
Alternative Catalog Number	BOT-BNUB2046-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant fragment (around aa 247-382) of human ZAP70 protein (exact sequence is proprietary)
Product Description	ZAP70 is a 70 kDa protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. ZAP70 protein is expressed in leukemic cells of approximately 25% of chronic lymphocytic leukemia (CLL) case...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[ZAP70/2046]
Molecular Weight	70 kDa

UniProt	P43403
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/m for coating order Ab without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user