

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

TACSTD2 / TROP2 (Epithelial Marker) (TACSTD2/2153), 0.2mg/mL, Clone: [TACSTD2/2153], Mouse, Monoclonal BOT-BNUB2153-100

Article Name	TACSTD2 / TROP2 (Epithelial Marker) (TACSTD2/2153), 0.2mg/mL, Clone: [TACSTD2/2153], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB2153-100
Supplier Catalog Number	BNUB2153-100
Alternative Catalog Number	BOT-BNUB2153-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant fragment of human TACSTD2 protein (around aa 31-274) (exact sequence is proprietary)
Product Description	TACSTD2 is a cell surface glycoprotein receptor. It is a single pass type I membrane protein containing one thyroglobulin type-1 domain, an epidermal growth factor-like repeat, a phosphatidylinositol binding site and tyrosine phosphorylation sites ne...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[TACSTD2/2153]

Molecular Weight	40 kDa
UniProt	P09758
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user